Rev. 11/98 . AGBX-2 CIP

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants

Michael Gallo et al.

Serial No.

09/375,924

Filed:

August 17, 1999

For

GENERATION OF MODIFIED MOLECULES WITH

INCREASED SERUM HALF-LIVES

Art Unit

1643

Hon. Assistant Commissioner for Patents Washington, D.C. 20231

RECEIVED

MAY 23 2000

TECH CENTER 1609/2800

TRANSMITTAL LETTER FOR INFORMATION DISCLOSURE STATEMENT

Sir:

Transmitted herewith is an Information Disclosure Statement in the above-identified application. This Statement is submitted more than three months from the application filing date and after the mailing date of the first Office Action on the merits, but before the mailing date of either a final action under 37 C.F.R. § 1.113, or a notice of allowance under 37 C.F.R. § 1.311.

In accordance with 37 C.F.R. § 1.97, this Statement is accompanied by:

- [] a certification as required under 37 C.F.R. § 1.97(e)(1) or § 1.97(e)(2);
- [X] the fee for \$240.00 as set forth in 37 C.F.R. \$1.17(p).

05/22/2000 MPRASASO 00000049 09375924

01 FC:126

240.00 DP

The Commissioner is hereby authorized to charge payment of any additional fees required in connection with the accompanying Information Disclosure Statement, or credit any overpayment, to Deposit Account No. 06-1075. A duplicate copy of this letter is transmitted herewith.

Respectfully submitted,

15 MAY 2020
Date

Daniel M. Becker Reg. No. 38,376

Attorney for Applicants

FISH & NEAVE 1251 Avenue of the Americas New York, N. Y. 10020 (650) 617-4000

I hereby Certify th Class Mail in an Er Addressed to: ASSISTANT COMMISSIONER FOR PATERTS WASHINGTON, D.G. 20231, ON

2000

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants : Michael Gallo et al.

Serial No. : 09/375,924

Filed: : August 17, 1999

For : GENERATION OF MODIFIED MOLECULES WITH

INCREASED SERUM HALF-LIVES

Art Unit : 1643

Hon. Assistant Commissioner for Patents

Washington, D.C. 20231

INFORMATION DISCLOSURE STATEMENT

KW 23 2007

Sir: TECH GENTER TOTAL T

Pursuant to 37 C.F.R. §§ 1.56, 1.97 and 1.98, applicants hereby make the following documents of record in the above identified application:

U.S. Patents

Axel et al.	4,399,216	08/1983
Kaufman	4,740,461	04/1998
Kaufman et al.	4,912,040	03/1990
Clark et al.	4,959,455	09/1990
Stec et al.	5,151,510	09/1992
Lonberg et al.	5,545,806	08/1996
Surani et al.	5,545,807	08/1996
Rostoker et al.	5,625,825	04/1997
Presta et al.	5,739,277	04/1998

Foreign Patent Documents

0 463 151	. B1	Europe	06/1996
0 770 628	A	Europe	05/1997
WO 91/082	98	WIPO	06/1991
WO 92/039	18	WIPO	03/1992
WO 92/226	345	WIPO	12/1992
WO 93/122	:27	WIPO	06/1993
WO 93/223	32	WIPO	11/1993
WO 94/026	502	WIPO	02/1994
WO 94/046	i89	WIPO	03/1994
WO 94/255	85	WIPO	11/1994
WO 96/085	512	WIPO	03/1996
WO 96/184	12	WIPO	06/1996
WO 96/324	78	WIPO	10/1996
WO 96/337	'35	WIPO	10/1996
WO 96/340	96	WIPO	10/1996
WO 97/346	31	WIPO	09/1997
WO 97/433	316	WIPO	11/1997
WO 97/443	362	WIPO	11/1997
WO 98/248	184	WIPO	06/1998
WO 98/248	193	WIPO	06/1998
WO 98/318		WIPO	07/1998

Other Publications

Anand et al., "Construction of yeast artificial chromosome libraries with large inserts using fractionation by pulsed-field gel electrophoresis." *Nucl. Acids Res.* 17:3425-3433 (1989)

Artandi et al. "Monoclonal IgM rheumatoid factors bind IgG at a discontinuous epitope comprised of amino acid loops from heavy-chain constant-region domains 2 and 3", *Proc Natl Acac Sci USA* 89:94-98 (1992)

Berman et al. "Content and organization of the human Ig VH locus: definition of three new V_H families and linkage to the Ig C_H locus." *EMBO J.* 7:727-738 (1988)

Bowie et al., "A method to identify protein sequences that fold into a known three-dimensional structure," *Science* 253:164 (1991)

Brambell et al., "A theoretical model of γ -Globulin Catabolism," Nature 203:1352-1355 (1964)

Brambell, "The transmission of immunity from mother to young and the catabolism of immunoglobulins," *The Lancet* II:1087-1093 (1966)

Brezinschek et al., "Analysis of the heavy chain repertoire of human peripheral B-cells using single-cell polymerase chain reaction." J. Immunol. 155:190-202 (1995)

Brownstein et al., "Isolation of single-copy human genes from a library of yeast artificial chromosome clones." *Science* 244:1348-1351 (1989)

Bruggeman et al., "A repertoire of monoclonal antibodies with human heavy chains from transgenic mice," *PNAS USA* 86:6709-6713 (1989)

Bruggemann et al., "Generation of antibody repertoires in transgenic mice," *Methods*: A Companion to Methods in Enzymology 2:159-165 (1991)

Bruggemann et al., "Human antibody production in transgenic mice: expression from 100 kb of the human IgH locus." Eur. J. Immunol. 21:1323-1326 (1991)

Bruggemann et al., "Strategies for expressing human antibody repertoires in transgenic mice." *Immunology Today*:17-391-397 (1996)

Burmeister et al., Crystal structure of the complex of rat neonatal Fc receptor with Fc, " Nature 372: 379-83 (1994)

Chen et al. "Immunoglobulin gene rearrangement in B-cell deficient mice generated by targeted deletion of the J_{H} locus" <code>International Immunology 5:647-656 (1993)</code>

Choi et al. "Transgenic mice containing a human heavy chain immunoglobulin gene fragment cloned in a yeast artificial chromosome" *Nature Genetics* 4:117-123 (1993)

Chothia & Lesk, "Canonical structures for the hypervariable regions of immunoglobulins," *J. Mol. Biol.* 196:901-917 (1987)

Chothia et al., "Conformations of immunoglobulin hypervariable regions," *Nature* 342:878-883 (1989)

Cook, G.P. and Tomlinson, I.M., "The human immunoglobulin $V_{\rm H}$ repertoire." Immunology Today 16:237-242 (1995)

Cox et al., "A directory of human germ-line Vx segments reveals a strong bias in their usage." Eur. J. Immunol. 24:827-836 (1994)

Dariavach et al., "The mouse IgH 3'-enhancer." Eur. J. Immunol. 21:1499-1504 (1991)

Den Dunnen et al., "Reconstruction of the 2.4 Mb human DMD-gene by homologous YAC recombination." Human MolecularGenetics 1:19-28 (1992)

Dima et al., "Effect of protein A and its fragment B on the catabolic and Fc receptor sites of IgG," Eur. J. Immunol. 13: 605 (1983)

Ellerson et al., "Structure and function of immunoglobulin domains," J. Immunol. 116:510 (1976)

Evans et al., "Design of nonpeptidal ligands for a peptide receptor: cholecystokinin antagonists," *J. Med. Chem.* 30:1229 (1987)

Fahey and Robinson, "Factors controlling serum γ -globulin concentration," A.G. J Exp. Med 118: 845-868 (1963)

Fauchere, "Elements for the rational design of peptide drugs," J. Adv. Drug Res. 15:29 (1986)

Feeney, A.J. "Lack of N regions in fetal and neonatal mouse immunoglobulin V-D-J junctional sequences." J. Exp. Med. 172:137-1390 (1990)

Fishwild et al., "High-avidity human IgGkmonoclonal antibodies from a novel strain of minilocus transgenic mice." *Nature Biotech*. 14:845-851 (1996)

Flanagan, J.G. and Rabbitts, T.H., "Arrangement of human immunoglobulin heavy chain constant region genes implies evolutionary duplication of a segment containing Y, ϵ , and α genes." Nature 300:709-713 (1982)

Fundamental Immunology Ch. 7 (Paul, W., ed., 2nd ed. Raven Press, N.Y. (1989)

Galfre, G. and Milstein, C., "Preparation of monoclonal antibodies: strategies and procedures." *Methods Enzymol*. 73:3-46 (1981)

Gemmill et al., "Protocols for pulsed field gel electrophoresis: Separation and detection of large DNA molecules." Advances in Genome Biology 1:217-251 (1992)

Ghetie et al., "Increasing the serum persistence of an IgG fragment by random mutagenesis," *Nature Biotechnology* 15:637 (1997)

Ghetie and Ward, "FcRn: the MHC class I-related receptor that is more than an IgG transporter," *Immunol Today* 18:592-598 (1997)

Gill et al., "Monoclonal anti-epidermal growth factor receptor antibodies which are inhibitors of epidermal growth factor binding and antagonists of epidermal growth factorstimulated tyrosine protein kinase activity." *J. Biol. Chem.* 259:7755 (1984)

Green et al., "Antigen-specific human monoclonal antibodies from mice engineered with human Ig heavy and light chain YACs." Nature Genetics 7:13-21 (1994)

Hermanson et al., "Rescue of end fragments of yeast artificial chromosomes by homologous recombination in yeast." *Nucleic Acids Res.* 19:4943-4948 (1991)

Huber et al., "The human immunoglobulin x locus. Characterization of the partially duplicated L regions." Eur. J. Immunol. 23:2860-2967 (1993)

Humphrey and Fahey, "The metabolism of normal plasma proteins and gamma-myeloma protein in mice bearing plasma-cell tumors," J. Clin. Invest. 40:1696-1705 (1961)

Immunoglobulin Genes pp. 259-274 (Honjo et al. eds., Academic Press Limited, San Diego, CA (1989)

Jakobovits, A., "Humanizing the mouse genome." Current Biology 4:761-763 (1994)

Jakobovits, A. et al., "Germ-line transmission and expression of a human-derived yeast artificial-chromosome." *Nature* 362:255-258 (1993)

Jakobovits, A. et al., "Analysis of homozygous mutant chimeric mice: Deletion of the immunoglobulin heavy-chain joining region blocks B-cell development and antibody production." *Proc. Natl. Acad. Sci. USA* 90:2551-2555 (1993)

Jakobovits, A., "Production of fully human antibodies by transgenic mice." Current Opinion in Biotechnology 6:561-566 (1995)

Junghans, "Finally! The brambell receptor (F_cR_B) " Immunologic Res. 16:29-57 (1997)

Junghans and Waldmann, "Metabolism of Tac (IL2Rα): physiology of cell surface shedding and renal catabolism, and suppression of catabolism by antibody binding," *J. Exp. Med* 183, 1587-1602 (1996)

Junghans et al., "The protection receptor for IgG catabolism is the β_2 -microglobulin-containing neonatal intestinal transport receptor," *Proc Natl Acac Sci USA* 93:5512-5516 (1996)

Kawamoto et al., "Growth stimulation of A431 cells by epidermal growth factor: Identification of high affinity receptors for EGF by an anti-receptor monoclonal antibody." *Proc. Nat. Acad. Sci.*, *USA* 80:1337-1341 (1983)

Kim et al., "Catabolism of the murine IgG1 molecule: evidence that both CH2-CH3 domain interfaces are required for persistence of IgG1 in the circulation of mice," *Scand J Immunol* 40:457-465 (1994)

Kim et al., "Evidence that the hinge region plays a role in maintaining serum levels of the murine IgG1 molecule," *Mol Immunol* 32:467-475 (1995)

Kim et al., "Localization of the site of the murine IgG1 molecule that is involved in binding to the murine intestinal Fc receptor," Eur J. Immunol. 24:2429-2434 (1994)

Kim et al., "Identifying amino acid residues that influence plasma clearance of murine IgG1 fragments by site-directed mutagenesis," Eur. J. Immunol. 24:542-548 (1994)

Knauf et al. "Relationship of Effective Molecular Size to Systemic Clearance in Rats of Recombinant Interleukin-2 Chemically Modified with Water-soluble Polymers," *J. Biochem.* 263:15064-15070 (1988)

Kostelny et al., "Formation of a bispecific antibody by the use of leucine zippers," J. Immunol. 148:1547-1553 (1992)

LaPlanche et al., "Phosphorothioate-modified oligodeoxyribonucleotides, III. NMR and UV spectroscopic studies of the Rp-Rp, Sp-Sp, and Rp-Sp duplexes, [d(GGsAATTCC)]₂, derived from diastereomeric O-ethyl phosphorothioates," Nucl. Acids Res. 14:9081 (1986)

Lonberg et al., "Antigen-specific human antibodies from mice comprising four distinct genetic modifications." *Nature* 368:856-859 (1994)

Lusti-Narasimhan et al., "Mutation of Leu 25 and Val 27 introduces CC chemokine activity into interleukin-8." *J. Biol. Chem.* 270:2716-2721 (1995)

Marks et al., "Oligonucleotide primers for polymerase chain reaction amplification of human immunoglobulin variable genes and design of family-specific oligonucleotide probes." *Eur. J. Immunol.* 21:985-991 (1991)

Mason and Williams, "The kinetics of antibody binding to membrane antigens in solution and at the cell surface," $Biochem\ J\ 187:1-20$ (1980)

Masson, "Elimination of infectious antigens and increase of IgG catabolism as possible modes of action of IVIg," J. Autoimmunity 6:683-689 (1993)

Matsuda et al., "Structure and physical map of 64 variable segments in the 3' 0.8-megabase region of the human immunoglobulin heavy-chain locus." Nature Genetics 3:88-94 (1993)

Max, E. Molecular genetics of immunoglobulins. Fundamental Immunology. 315-382 (Paul, WE, ed., New York: Raven Press (1993)

McFarlane, "The behavior of I¹³¹-labeled plasma proteins In Vivo," Ann NY Acad Sci 70:19-25 (1957)

Medesan et al., "Delineation of the amino acid residues involved in transcytosis and catabolism of mouse IgG1¹," *J Immunol* 158:2211-2217 (1997)

Medesan et al., "Localization of the site of the IgG molecule that regulates maternofetal transmission in mice," $Eur.\ J.\ Immunol.\ 26:2533-2536\ (1996)$

Mendez et al., "Analysis of the structural integrity of YACs comprising human immunoglobulin genes in yeast and in embryonic stem cells." *Genomics* 26:294-307 (1995)

Mendez et al., "A set of YAC targeting vectors for the interconversion of centric and acentric arms." Cold Spring Harbor Laboratory Press, Genome Mapping and Sequencing meeting, 163 (1993)

Mendez et al., "Functional transplant of megabase human immunoglobulin loci recapitulates human antibody response in mice", *Nature Genetics* 15:146-156 (1997)

Needleman and Wunsch, "A general method applicable to the search for similarities in the amino acid sequence of two proteins," J. $Mol.\ Biol.\ 48:443\ (1970)$

Nose and Wigzell, "Biological significance of carbohydrate chains on monoclonal antibodies," *Proc. Natl. Acad. Sci. USA* 80:6632 (1983)

Pearson and Lipman, "Improved tools for biological sequence comparison," Proc. Natl. Acad. Sci. (U.S.A.) 85:2444 (1988)

Pollock et al., "Intravascular metabolism of normal and mutant mouse immunoglobulin molecules," *Eur. J. Immunol.* 20:2021-2027 (1990)

Proc. Natl. Acad. Sci., "Construction and characterization of a yeast artificial chromosome," 87:4256 (1990)

Raghavan et al., "Investigation of the interaction between the class I MHC-related Fc receptor and its immunoglobulin G ligand," Immunity 1:303-315 (1994)

Raghavan et al., "Effects of receptor dimerization on the interaction between the class I major histocompatibility complex-related Fc receptor and IgG," *Proc Natl Acac Sci USA* 92:11200-11204 (1995)

Raghavan et al., "Analysis of the pH dependence of the neonatal Fc receptor/immunoglobulin G interaction using antibody and receptor variants," *Biochemistry* 34:14649-14657 (1995)

Ray, S. and Diamond, B., "Generation of a fusion partner to sample the repertoire of Splenic B-cells destined for apoptosis." *Proc. Natl. Acad. Sci. USA* 91:5548-5551 (1994)

Rizo and Gierasch, "Constrained peptides: models of bioactive peptides and protein substructures," Ann. Rev. Biochem. 61:387 (1992)

Sambrook et al. *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)

Sato et al., "Biological effects in vitro of monoclonal antibodies to human epidermal growth factor receptors" Mol. Biol. Med. 1:511-529 (1983)

Schiestl, R.H. and Gietz, R.D., "High efficiency transformation of intact yeast cells using stranded nucleic acids as a carrier." Curr. Genet. 16:339-346 (1989)

Segal et al., "The role of non-immune IgG in controlling IgG-mediated effector functions," Mol Immunol 20:1177-1189 (1983)

Sell and Fahey, "Relationship between γ -globulin metabolism and low serum γ -globulin in germfree mice," *J Immunol* 93:81-87 (1964)

Sell, "Evidence for species' differences in the effect of serum γ -globulin concentration on γ -globulin catabolism," *J. Exp. Med.* 120:967-986 (1964)

Silverman et al., "Meiotic recombination between yeast artificial chromosomes yields a single clone containing the entire BCL2 protooncogene." *Proc. Natl. Acad. Sci. USA* 87:9913-9917 (1990)

Smith and Waterman, "Comparison of biosequences," Adv. Appl. Math. 2:482 (1981)

Songsivilai & Lachmann, "Bispecific antibody: a tool for diagnosis and treatment of disease," *Clin. Exp. Immunol.* 79:315-321 (1990)

Spiegelberg in 19 Advances in Immunology F. J. Dixon and H. G. Kinkel, eds. 259-294 (Academic Press, NY: 1974)

Spiegelberg and Wiegle, "The catabolism of homologous and heterologous 7S gamma globulin fragments," *J. Exp. Med.* 121:323-338 (1965)

Srivastava, A. and Schlessinger, D., "Vectors for inserting selectable markers in vector arms and human DNA inserts of yeast artificial chromosomes (YACs)." Gene 103:53-59 (1991)

Stec et al., "Automated solid-phase synthesis, separation, and stereochemistry of phosphorothioate analogues of oligodeoxy-ribonucleotides," J. Am. Chem. Soc. 106:6077 (1984)

Stein et al., "Physicochemical properties of phosphorothioate oligodeoxynucleotides," *Nucl. Acids Res.* 16:3209 (1988)

Tao and Morrison, "Role of carbohydrate in the structure and effector functions mediated by the human IgG constant region," $J.\ Immunol.\ 143:2595\ (1989)$

Taylor et al., "A transgenic mouse that expresses a diversity of human sequence heavy and light chain immunoglobulins." *Nucleic Acids Research* 20:6287-6295 (1992)

Taylor et al., "Human immunoglobulin transgenes undergo rearrangement, somatic mutation and class switching in mice that lack endogenous IgM." *International Irnmunology* 6:579-591 (1994)

Thornton et al., "Prediction of progress at last," *Nature* 354:105 (1991)

Tuaillon et al. "Analysis of direct and inverted DJ_H rearrangements in a human Ig heavy chain transgenic minilocus" J. Immunol. 154:6453-6465 (1995)

Tuaillon et al., "Human immunoglobulin heavy-chain minilocus recombination in transgenic mice: gene-segment use in m and Y transcripts." *Proc. Natl. Acad. Sci. USA* 90:3720-3724 (1993)

Uhlmann and Peyman, "Antisense Oligonucleotides: A new therapeutic principle," Chemical Reviews 90:543 (1990)

Vaughan et al., "Human antibodies with subnanomolar affinities isolated from a large non-irnmunized phage display library."

Nature Biotech. 14:309-314 (1996)

Vaughn and Bjorkman, "Structural basis of pH-dependent antibody binding by the neonatal Fc receptor," Research Article, 63-73, 1998

Vaughn and Bjorkman, "High-affinity binding of the neonatal Fc receptor to its IgG ligand requires receptor immobilization," Biochem, 36:9374-9380 (1997)

Veber and Freidinger, "The design of metabolically-stable peptide analogs," Trends In NeuroScience p.392-396 (1985)

Wagner et al., "The diversity of antigen-specific monoclonal antibodies from transgenic mice bearing human immunoglobulin gene miniloci." Eur. J. Irnmunol. 24:2672-2681 (1994).

Waldman and Strober, "Metabolism of immunoglobulins," Progress in Allergy 13: 1-110, (1969)

Waldmann and Ghetie "Catabolism of Immunoglobulins," *Progress in Immunol*. 1:1187-1191 (Academic Press, New York: 1971)

Waldmann and Jones, "The role of cell-surface receptors in the transport and catabolism of immuno-globulins," *Protein Turnover* 9:5-23 (1973)

Wallace and Rees, "Studies on the immunoglobulin-G Fc-Fragment receptor from neonatal rat small intestine," *Biochem J* 188: 9-16 (1980)

Wawrzynczak et al., "Recombinant mouse monoclonal antibodies with single amino acid substitutions affecting Clq and high affinity Fc receptor binding have identical serum half-lives in the balb/c mouse," *Molec. Immunol.* 29:221-227 (1992)

Wawrzynczak et al., "Blood clearance in the rat of a recombinant mouse monoclonal antibody lacking the N-linked oligosaccharide side chains of the C_{H2} domains," $Mol.\ Immunol.\ 29:213-220\ (1992)$

Weichhold et al., "The human immunoglobulin k locus consists of two copies that are organized in opposite polarity." *Genomics* 16:503-511 (1993)

Wochner et al., "The role of the kidney in the catabolism of Bence Jones proteins and immunoglobulin fragments," *J. Exp. Med.* 126:207 (1967)

Yamada, M. et al., "Preferential utilization of specific immunoglobulin heavy chain diversity and joining segments in adult human peripheral blood B lymphocytes." J. Exp. Med. 173:395-407 (1991)

Yasmeen et al., "The structure and function of immunoglobulin domains," J. Immunol. 116:518 (1976)

Zon et al. Oligonucleotides and Analogues: A Practical Approach, pp. 87-108 (F. Eckstein, Ed., Oxford University Press, Oxford England (1991)

Zon et al., "Phosphorothioate oligonucleotides: chemistry, purification, analysis, scale-up and future directions," Anti-Cancer Drug Design 6:539 (1991)

Zuckier et al., "The use of severe combined immunodeficiency mice to study the metabolism of human immunoglobulin G," *Cancer* 73:794-799 (1994)

Zuckier et al., "Immunologic and pharmacologic concepts of monoclonal antibodies," Semin. Nucl. Med. 19:166-186 (1989)

Copies of the aforementioned documents, which are listed on the accompanying Form PTO-1449 (submitted in duplicate), are enclosed herewith.

It is respectfully requested that these documents be

(1) fully considered by the Patent and Trademark Office during
the examination of this application; and (2) printed on any
patent which may issue on this application. Applicants request
that a copy of Form PTO-1449 (submitted in duplicate herewith),
as considered and initialed by the Examiner, be returned with the
next communication.

This information disclosure statement is understood to complement the Examiner's own search, and shall not be construed as a representation that a search has been made. The filing of this information disclosure statement shall not be construed to be an admission that the information cited in the statement is, or is considered to be, material to patentability as defined in § 1.56. Applicant reserves the right to challenge the applicability as prior art of any of the above-cited references.

An early and favorable action is respectfully requested.

14 MAY 2009
Date

Daniel M. Becker Reg. No. 38,376

Respectfully

-Attorney for Applicants

thereby Cordly that the
Correspondence is being
Deposited with the U.S.
Postal Service as First
Class Mail in an Envelope
Addressed to: ASSISTANT
COMMISSIONER FOR PATENTS
WASHINGTON, D.C. 20231, CNI

Signature of Person Signing

FISH & NEAVE 1251 Avenue of the Americas New York, N. Y. 10020 (650) 617-4000